

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Statistical reports of States and cities of the United States—Yearly and monthly.

MICHIGAN.—Reports to the State board of health, Lansing, for the week ended March 16, 1901, from 66 observers, indicate that diarrhea, enteric fever, pleuritis, pneumonia, and whooping cough were more prevalent, and inflammation of kidneys less prevalent than in the preceding week. Cerebro-spinal meningitis was reported present at 4, whooping cough at 13, diphtheria at 27, measles at 28, enteric fever at 40, scarlet fever at 103, phthisis pulmonalis at 172, and smallpox at 75 places.

Reports to the State board of health, Lansing, for the week ended March 23, 1901, from 84 observers, indicate that diphtheria, enteric fever, pleuritis, and pneumonia were less prevalent than in the preceding week. Cerebro-spinal meningitis was reported present at 4, whooping cough at 10, measles at 25, diphtheria at 26, enteric fever at 41, scarlet fever at 96, phthisis pulmonalis at 182, and smallpox at 77 places.

NEW JERSEY—Paterson.—Month of February, 1901. Estimated population, 106,515. Total number of deaths, 154, including diphtheria, 4; enteric fever, 1, and 18 from phthisis pulmonalis.

NEW YORK—Rochester.—Month of February, 1901. Census population, 162,608. Total number of deaths, 222, including diphtheria, 2' and 19 from tuberculosis.

UTAH—Salt Lake City.—Month of January, 1901. Census population, 53,531. Total number of deaths, 70, including diphtheria, 5; la grippe, 7, and 5 from tuberculosis. Month of February, 1901. Total number of deaths, 54, including diphtheria, 3; smallpox, 1, and 2 from tuberculosis.

WISCONSIN—*Milwaukee*.—Month of February, 1901. Census population, 285,315. Total number of deaths, 317, including diphtheria, 5; enteric fever, 3; measles, 2; scarlet fever, 2; whooping cough, 5, and 32 from tuberculosis.

Report of immigration at Boston for the week ended March 30, 1901.

## OFFICE OF U. S. COMMISSIONER OF IMMIGRATION, Port of Boston, March 31, 1901.

Number of alien immigrants who arrived at this port during the week ended March 30, 1901; also names of vessels and ports from which they came.

Date.	Vessel.	Where from.	No. of in migrants.
Mar. 24 Do Mar. 26 Mar. 28 Do Do Mar. 29	Steamship Boston Steamship Devouian Steamship Prince George Steamship Boston Steamship Halifax	Yarmouth, Nova Scotia	135 7 92 116 45
	Total		565